

## Thermophiles and Their Industrial Application: A Review

Ayisa T.T.<sup>1</sup>, Oyedokun N.O.<sup>2</sup>, Ideh R.<sup>2</sup>, Itoroabasi J.O.<sup>2</sup>, Ideh R.R.<sup>2</sup>, Egbi T.K.<sup>2</sup>, Ahmadu J.H.<sup>2</sup> and Orji S.L.<sup>2</sup>

<sup>1</sup>Biological Science Department, The Federal Polytechnic Bida, P.M.B. 55, Bida, Niger State, Nigeria.

<sup>2</sup>National Biotechnology Development Agency, Abuja, Nigeria.

Corresponding author: [ayisatimothy4u@gmail.com](mailto:ayisatimothy4u@gmail.com)

### Abstract

Thermophilic micro organisms are of important economic value due to their ability to produce thermostable extra cellular enzymes which have important biotechnological and industrial applications. It is well known that thermophilic activities are generally associated with protein stability. Thus, the proteins which are produced by these thermophilic micro organisms are in most cases more stable compared to their mesophilic counterparts. The mechanism of thermostability of the proteins that results in the molecular rigidity is related to a number of salt bridges, hydrogen and disulphide bonds and the packing of external residues. The biotechnological advantage of these thermostable enzymes is their ability to withstand elevated temperatures, reducing risk of contamination by mesophiles. Several cellulases, xylanases and pectinases have been applied in industrial and biotechnological processes such as oil recovery, production of animal feed, biobleaching of paper pulp, obtaining biofuel from cellulosic wastes just to mention a few. This review captures thermophilic micro organisms and their thermostable enzymes in a number of industrial and biotechnological applications.

**Keywords:** Thermophiles, Biotechnology Industrial, Enzymes Thermostable.

### Introduction

Organisms with an optimum temperature for growth between 60 and 80°C are generally designated as thermophiles, while those growing optimally above 80°C are referred to as hyper thermophiles (Santos and Da Costa, 2016). Thermophilic bacteria are microbes that mostly inhabit hot springs, live and survive in temperatures above 70°C. As a consequence of growth at high temperatures and unique macromolecular properties, thermophiles can possess high metabolism, physically and chemically stable enzymes and lower growth but higher end product yields than similar mesophilic species (Sprott *et al.*, 2018).

Natural environments for anaerobic thermophiles range from terrestrial volcanic sites (including solfatara fields) with temperatures slightly above ambient temperature, to submarine hydrothermal systems (sediments, submarine volcanoes, fumaroles and vents) with temperatures exceeding 300 °C, subterranean sites such as oil reservoirs, and solar heated surface soils with temperatures up to 65 °C. There are also human-made hot environments such as compost piles (usually around 60–70 °C but as high as 100°C) slag heaps, industrial processes and water heaters (Oshima and Moriya, 2019).

The ubiquitous nature of the thermophiles is attested to by the great variety of sources from which they have been isolated from freshly fallen snow to the sands of the Sahara Desert. They have been found to occur in the air, the soil of temperate and tropical regions, salt and fresh water, both cold and thermal (Golikova, 2018).

Factors affecting heat tolerance of thermophilic organisms are as follows:

- i. **Permeability:** cell membranes effectively function as a permeability barrier, controlling the in-flow and out-flow of low-molecular weight compounds. The permeability of fatty acyl ester lipid membranes is highly temperature dependent and their phase-transition temperature is dependent on the fatty acid composition, so when the growth temperature shifts, the fatty acid composition of membrane lipids is quickly regulated (Koga, 2019).
- ii. **Chemical stability:** thermophilic organisms are able to grow at high temperature due to the chemical stability of their membrane lipids (Koga, 2019).
- iii. **Temperature:** lipids that increase in proportion to an increase in growth temperature may be designated as “thermophilic lipids.” In the extremely thermophilic environment, methanoarchaea *Methanocaldococcus jannaschii* have been reported. When the growth temperature increases from 45 to 65 °C, the diether lipids (archaeol-based lipids) decrease from 80 to 20 %, while the standard caldarchaeol-based and cyclic archaeol-based lipids increase from 10 to 40 %, respectively (Sprott *et al.*, 2018).
- iv. **G+C content:** rRNA and tRNA molecules of thermophilic bacteria have higher G+C contents than mesophiles (Galtier and Lobry, 2018). Because the GC base pair forms more hydrogen bonds than the AT base pair, higher GC contents in the double-stranded stem region improves thermostability of the RNA molecules (Lao and Forsdyke, 2018).
- v. **Proteins:** the surface regions of thermophilic proteins have fewer (non-charged) polar amino acids and more charged amino acids, and these charged residues result in an increased number of intramolecular salt bridges (Thompson and Eisenberg, 2018).

### Industrial Applications of Thermophiles

Thermophiles have shown tremendous promise in terms of their applications in modern biotechnology. Some of the high end applications of these thermophiles have been elucidated below:

#### Bioconversion of Lignocellulose to Hydrogen

Although many reported microorganisms possess the capability of cellulose hydrolysis or hydrogen production ( $H_2$ ), no conclusive research has been able to clarify that both of these capabilities are possessed in a single microorganism. Hot springs are a potential source for thermophilic hydrogen ( $H_2$ ) and ethanol producing microorganisms. Compared with mesophiles, thermophiles are thought to be more robust for cellulose degradation and hydrogen production. In particular, the rate of cellulolysis is presumably more rapid at elevated temperatures (Wiegel and Ljungdahl 2019). As a result, thermophilic microorganisms isolated from various environments are an attractive prospect for cellulolytic biohydrogen production (CBP) from complex lignocellulosic biomass. The co-cultures of thermophilic cellulolytic bacterium *Clostridium thermocellum* with non-cellulolytic thermophilic anaerobic bacteria and the extremely thermophilic cellulolytic bacterium *Caldicellulosiruptor saccharolyticus* have been used for CBP-based hydrogen production (Liu *et al.*, 2019). Several species of genus *Thermoanaerobacterium* including *T. thermosaccharolyticum*, *T. polysaccharolyticum*, *T. zeae*, *T. lactoethylicum*, *T. aotearoense*, and *T. saccharolyticum* possess the capability to utilize various macromolecules accompanied by  $H_2$  production (Ganghofer *et al.*, 2019).

Several anaerobic thermophiles have been shown to utilize cellulose, including *Clostridium thermocellum*, *Clostridium straminisolvans*, *Clostridium stercorarium*, *Caldicellulosiruptor saccharolyticus*, and *Caldicellulosiruptor obsidiansis* (Rainey *et al.*, 2019).

Hydrogen may be the fuel of the future once hydrogen fuel cells for propelling cars are perfected (McAlister, 2019). When oxygen and hydrogen are combined in a fuel cell, they provide electricity and a little heat, giving water as the only waste product. The hydrogen car will be clean because it will not discharge nitrogen oxides and carbon dioxide. Hydrogen can be obtained cheaply using special enzymes (extremozymes) by the transformation of cellulose into glucose sugar, then converting the glucose product and its byproduct, gluconic acid into hydrogen (Woodward *et al.*, 2018).

The extremely thermophilic bacterium *C. owensensis* has comprehensive hemicellulase and cellulase system. The enzymes of *C. owensensis* had high ability for degrading the hemicellulose of native lignocellulosic biomass. High temperature pre-hydrolysis on native lignocellulosic biomass by the extra-enzyme of *C. owensensis* could greatly improve the glucan conversion rate, making almost the same contribution as steam-exploded pretreatment (Peng *et al.*, 2019).

### Conversion of Glycerol to Lactate

Bioprospecting efforts for exploring novel biocatalytic molecules with unique properties have inspired the design and construction of a wider variety of artificial metabolic pathways (Bond-Watts *et al.*, 2018). Employment of enzymes derived from thermophiles and hyperthermophiles enables the simple preparation of catalytic modules with excellent selectivity and thermal stability (Ye *et al.*, 2019). These excellent stabilities of thermophilic enzymes allow greater flexibility in the operational conditions of in vitro bioconversion systems.

Generally, 10 kg of crude glycerol, which is the primary byproduct of the biodiesel industry, is released for every 100 kg of biodiesel and the growing production of biodiesel has resulted in a worldwide surplus of crude glycerol (Nguyen *et al.*, 2018). An artificial in vitro metabolic pathway for the conversion of glycerol to lactate has been constructed. The in vitro pathway consisted of nine thermophilic and hyperthermophilic enzymes and designed to balance the intrapathway consumption and regeneration of cofactors (Okano *et al.*, 2019).

### Conversion of D-Xylose into Ethanol

Thermophilic anaerobic bacteria could be promising candidates for conversion of hemicellulose or its monomers (xylose, arabinose, mannose and galactose) into ethanol with a satisfactory yield and productivity. A number of thermophilic enrichment cultures, and new isolates of thermophilic anaerobic bacterial strains growing optimally at 70–80 °C for their ethanol production from D-xylose have been isolated from hot springs, paper pulp mills and brewery waste water. The species investigated so far are *Thermoanaerobacter ethanolicus* (Kannan and Mutharasan 2020), *Clostridium thermocellum* (Viljoen *et al.*, 2018), *Clostridium thermohydrosulfuricum* (Cook and Morgan 2019,) (reclassified as *Thermoanaerobacter thermohydrosulfuricus*) *Thermoanaerobium Brockii* (reclassified as *Thermoanaerobacter Brockii*) (Lee *et al.*, 2019), *Clostridium thermosaccharolyticum* (reclassified as *Thermoanaerobacterium thermosaccharolyticum*) (Collins *et al.*, 2019) and *Thermoanaerobacterium saccharolyticum* B6A (Weimer, 2020). To obtain a viable bio-ethanol production, all the carbohydrate constituents of lignocellulosic biomass need to be converted into ethanol (Von Sivers and Zacchi, 2020). Xylan is mainly composed of D-xylose and it has been reported that most of the isolates produced ethanol as the main end fermentation product from both xylose and xylan (Sommer *et al.*, 2016). Among the advantages, thermophiles have broad substrate spectra and can degrade both hexoses and pentoses simultaneously; some thermophiles natively degrade complex carbohydrates;

they operate at temperatures that minimize contamination risk of mesophiles. Additionally, recent advances have improved ethanol yields by using genetic engineering, often by knocking out metabolic pathways to other end products. This has minimized the perceived advantage of mesophiles over thermophiles considerably, although no large scale bioethanol plants with genetically modified microbes are currently operating (Scully and Orlygsson, 2019).

#### **Biodegradation of Petroleum Hydrocarbons**

Thermophiles have also been utilized for the microbial degradation of crude oil and refined petroleum pollutants. Foght and McFarlane (2018) studied the growth of extremophiles on petroleum hydrocarbon. Some potential applications are related to molecular genetics of polycyclic aromatic hydrocarbon degradation by bacteria. Also, the factors that control degradation and methods to enhance the ability of bacteria to degrade such pollutants in the environment have been studied. April *et al.*, (2018) studied the process of crude oil degradation by mixed populations, pure cultures, and genetic mutants. They demonstrated the loss of parent compounds and analyzed the products of bacterial attack on crude oils using gas chromatography, mass spectrometry. The use of thermophiles for biodegradation of hydrocarbons with low water solubility is of interest, as solubility and bioavailability, are enhanced at elevated temperatures. Thermophiles, predominantly bacilli, possess a significant potential for the degradation of environmental pollutants, including all major classes. Indigenous thermophilic hydrocarbon degraders are of special significance for the bioremediation of oil-polluted desert soil (Margesin and Schinner, 2019).

#### **Recovery of Heavy Metals**

As a result of increasing industrial activities, heavy metal contamination is a problem. Microorganisms can interact with heavy metals in a variety of ways that result in decreased metal mobility and solubility. The metal and sulfate-reducing bacteria have suitable physiology for metal precipitation and immobilization. The activities of these microbes provide metabolic products such as iron and hydrogen sulphide, which lead to mineral formation. These minerals can react with heavy metals, resulting in precipitation and hence detoxification (Chalaal and Islam, 2019). In order to understand the removal of such types of toxins, Chalaal and Islam (2019) used two strains of thermophilic bacteria belonging to the *Bacillus* family, isolated from hot water stream, to remove strontium from aqueous stream systems. These bacteria were able to concentrate strontium in one side of a two-compartment bioreactor. Immobilization of heavy metals using sulphide-producing microorganisms has been reported as an effective means of treating some metal-contaminated sites (Crawford and Crawford, 2019).

#### **Remediation of Textile Dyes**

Laccase enzyme purified from thermophile, *Geobacillus thermocatenulatus* MS5 is of very higher catalytic activity and are economic, highly stable at different temperatures and pH levels and can be used widely and effectively in the removal of the dyes that cause environmental pollution. Verma and Shirkot investigated the purified laccase enzyme for the removal of some dyes used in industry i.e., Remazole Brilliant Blue R (RBBR), Indigo carmine, Congo red, Brilliant green and Bromophenol blue. In case of Indigo carmine and congo red dye, 99 % of decolorization occurred after 48 h of incubation, followed by RBBR dye, Bromophenol Blue and Brilliant Green i.e., 98, 70 and 60 % respectively (Verma and Shirkot, 2018). Thermophilic lignolytic fungal cultures were isolated from soil/digested slurry/plant debris and were subjected for acclimatization to Remazol Brilliant Blue

(RBB) at 0.05 % concentration, in the malt extract broth (MEB). The results suggested the isolates as a useful tool for degradation of reactive dyes (Sahni and Gupta, 2018).

### Saccharification of Agricultural Residues

*Sporotrichum thermophile* LAR5 is an excellent fungal isolate having an ability to utilize crude agriculture-based materials as carbon and nitrogen sources to produce significant cellulose titre. Cellulase possesses desirable properties from industrial application point of view such as activity and stability over broad pH range and high temperatures and good saccharification ability on acid pretreated rice straw. It has been reported that considerable sugars are produced by enzymatic hydrolysis of acid-pretreated solids (3.5, 5.7, 7.9, 7.7 micromoles/ml from 1, 3, 5 and 7 % acid-pretreated solids, respectively) using the *S. thermophile* LAR5 cellulase (Bajaj *et al.*, 2018). Recombinant *S. thermophile* cellulase shows potential to hydrolyze variety of cellulosic substrates with a peculiarity that presence of lignin in various substrates enhances the degree of saccharification (Dimarogona *et al.*, 2019).

### Thermophilic Bacilli in Dairy Processing

Thermophilic bacilli are used as hygiene indicators of processed product, within the dairy processing context. This is because of the ability of these strains to form endospores and biofilms. The thermophilic bacilli, such as *Anoxybacillus flavithermus* and *Geobacillus spp.*, are an important group of contaminants in the dairy industry. Although these bacilli are generally not pathogenic, their presence in dairy products is an indicator of poor hygiene and high numbers are unacceptable to customers. In addition, their growth may result in milk product defects caused by the production of acids or enzymes, potentially leading to off-flavors (Burgess *et al.*, 2019). Many strains of genera *Lactobacillus* and *Bifidobacterium*, as well as some enterococci and yeasts, have been shown to possess probiotic properties with potential for prophylaxis and treatment of a range of gastrointestinal disorders (Varankovich *et al.*, 2019).

### Keratin Degradation

A novel thermophilic bacterium, *Fervidobacterium pennavorans*, belonging to the *Thermotogales* order, isolated from hot springs of Azores Island, grows optimally at 70 °C and pH 6.5. It is the first known thermophile that is able to degrade native feathers at high temperatures. With the help of these enzymes, feathers could be converted to defined products such as the rare amino acids, serine, cysteine and proline (Friedrich and Antranikian, 2019).

### Cancer Treatment

Asperjinone, a nor-neolignan, and Terrein, a suppressor of ABCG2-expressing breast cancer cells were isolated from thermophile *Aspergillus terreus*, which can restore drug sensitivity and could be the key to improve breast cancer therapeutics. Terrein, displayed strong cytotoxicity against breast cancer MCF-7 cells. Treatment with terrein significantly suppressed growth of ABCG2-expressing breast cancer cells. This suppressive effect was achieved by inducing apoptosis via activating the caspase-7 pathway and inhibiting the Akt signaling pathway, which led to a decrease in ABCG2-expressing cells and a reduction in the side-population phenotype (Liao *et al.*, 2019). Conventional chemotherapeutic agents are usually nonspecific towards cancerous cells and inhibit the progression of any dividing cells. The therapeutic potential of antitumor drugs is seriously limited by the

manifestation of serious side effects and drug resistance. So there is a need of agents that are more effective, more selective and may not cause drug resistance. According to Patent no. WO 2006/053445 A1, an invention is disclosed, whereby a composition of Bacteriocins derived from lactic acid bacteria and a carrier can be used for inhibiting proliferation of cancerous cells (Mehta *et al.*, 2018).

### Conclusion

The increasing number of patents indicates that there is a growing interest in the commercial applications of thermophiles. The demand for thermostable enzymes has increased tremendously in the past few years. Since only a very few species from this group of microorganisms have been isolated till date, there seems to be a large number of hyperthermophilic catalysts with unique properties awaiting discovery. These promising results can be exploited further for production of biotechnological important and industrially thermostable enzymes. This study widens the opportunities for further research to be conducted to explore more the immense significance of these strains especially *Thermomonas* isolates, where there is lack of intensive studies regarding this organism.

### Recommendations

- Government should encourage the use of Thermophiles in hospitals as it has a positive impact on the treatment of Cancer, in order to increase the chances of the survival of Cancer patients.
- Further research should be carried out on mechanism of action thermophiles; this is to enlighten the populace especially Food and Pharmaceutical industries of the dos and don'ts involving the use of thermophiles, so that it does not cause antagonistic effects.

### References

- April, T. M., Foght, J. M. and Currah, R. S. (2018). Hydrocarbon-degrading fungi isolated from flare pit soils in northern and western Canada. *Can. J. Microbiol.*, 46: 38-49.
- Bajaj, B. K., Sharma, M. and Rao, R. S. (2018). Agricultural residues for production of cellulase from *Sporotrichum thermophile* LAR5 and its application for saccharification of rice straw. *J. Mater. Environ. Sci.*, 5(5): 1454-1460.
- Bond-Watts, B. B., Bellerose, R. J. and Chang M. C. Y. (2011). Enzyme mechanism as a kinetic control element for designing synthetic biofuel pathways. *Nat. Chem. Biol.*, 7: 222-227.
- Burgess, S. A., Lindsay, D. and Flint, S. H. (2019). Thermophilic bacilli and their importance in dairy processing. *Int. J. Food Microbiol.*, 144: 215-225.
- Chalaal, O. and Islam, M. R. (2019). Integrated management of radioactive strontium contamination in aqueous stream systems. *J. Environ. Manage.* 61: 51-59.
- Collins, M. D., Lawson, P. A., Willems, A., Cordoba J. J., Fernandez- Garayzabal, J., Garcia, P., Cai, J., Hippe, H. and Farrow, A. E. (2019). The phylogeny of the genus *Clostridium*: proposal of five new genera and eleven new species combinations. *Int. J. Syst. Bacteriol.* 44: 812-826.
- Cook, G. M. and Morgan, H. W. (2019). Hyperbolic growth of *Thermoanaerobacter thermohydrosulfuricus* (*Clostridium thermohydrosulfuricus*) increases ethanol production in pH controlled batch culture. *Appl. Microbiol. Biotechnol.* 41: 84-89.
- Crawford, R. L. and Crawford, D. L. (2019). Bioremediation: principles and applications. *Appl. Microbiol. Biotechnol.*, 8: 220-300.
- Dimarogona, M., Topakas, E., Olsson, L. and Christakopoulos, P. (2019). Lignin boosts the cellulose performance of a GH-61 enzyme from *Sporotrichum thermophile*. *Bioresour. Technol.*, 110: 480-487.

- Foght, J. M. and McFarlane, D. M. (2018). Growth of extremophiles on petroleum. In: Seckbach J (ed) Enigmatic microorganisms and life in extreme environments. *Kluwer. Academic Publishers*, 1: 527-538.
- Friedrich, A. B. and Antranikian, G. (2019). Keratin degradation by *Fervidobacterium pennavorans*, a novel thermophilic anaerobic species of the order Thermotogales. *Appl. Environ. Microbiol.*, 62(8): 2875-2882.
- Galtier, N. and Lobry, J. R. (2018). Relationships between genomic G+C content, RNA secondary structures, and optimal growth temperature in prokaryotes. *J. Mol. Evol.*, 44(6): 632-636.
- Ganghofer, D., Kellermann, J., Staudenbauer, W. L. and Bronnenmeier, K. (2019). Purification and properties of an amylopullulanase, a glucoamylase, and an alphaglucohydrolase in the amylolytic enzyme system of *Thermoanaerobacterium thermosaccharolyticum*. *Biosci. Biotech. Bioch.* 62: 302-308.
- Golikova, S. M. (2018). Developments in industrially important Thermostable enzymes. *Bioresour. Technol.*, 89: 17-34.
- Kannan, V. and Mutharasan, R. (2020). Ethanol fermentation characteristics of *Thermoanaerobacter ethanolicus*. *Enzyme Microb. Technol.*, 7: 87-89.
- Koga, Y. (2019). Thermal adaptation of the archaeal and bacterial lipid membranes: Fermentation of moderately thermophilic bacilli on sucrose. *Archaea.*, 8: 663-954.
- Lao, P. J. and Forsdyke, D. R. (2018). Thermophilic bacteria strictly obey Szybalski's transcription direction rule and politely purine-load RNAs with both adenine and guanine. *Genome Res.*, 10(2): 228-236.
- Lee, Y., Jain, M. K., Lee, C., Lowe, S. E. and Zeikus, J. G. (2019). Taxonomic distinction of saccharolytic thermophilic anaerobes: description of *Thermoanaerobacterium xylanolyticum* gen nov, sp nov, and *Thermoanaerobacterium saccharolyticum* gen nov, sp nov; reclassification of *Thermoanaerobium brockii*, *Clostridium thermosulfurigenes* and *Clostridium thermohydrosulfuricum* E100-69 as *Thermoanaerobacter brockii* comb nov, *Thermoanaerobacterium thermosulfurigenes* comb nov, and *Thermoanaerobacterthermohydrosulfuricus* comb nov, respectively; and transfer of *Clostridium thermohydrosulfuricum* 39E to *Thermoanaerobacter ethanolicus*. *Int. J. Syst. Bacteriol.*, 43: 41-51.
- Liao, W. Y., Shen, C. N., Lin, L. H., Yang, Y. L., Han, H. Y., Chen, J. W., Kuo, S. C., Wu, S. H. and Liaw, C. C. (2019). Asperjinone, a nor-neolignan, and terrein, a suppressor of ABCG2-expressing breast cancer cells, from thermophilic *Aspergillus terreus*. *J. Nat. Prod.*, 75(4): 630-635.
- Liu, Y., Yu, P., Song, X. and Qu, Y. B. (2019). Hydrogen production from cellulose by coculture of *Clostridium thermocellum* JN4 and *Thermoanaerobacterium thermosaccharolyticum* GD17. *Int. J. Hydrogen Energ.*, 33: 2927-2933.
- Margesin, R. and Schinner, F. (2019). Biodegradation and bioremediation of hydrocarbons in extreme environments. *Appl. Microbiol. Biotechnol.*, 56: 650-663.
- Mehta, R., Arya, R., Goyal, K., Singh, M. and Sharma, A. K. (2018). Biopreservative and therapeutic potential of pediocin: Recent trends and future perspectives. *Recent Pat. Biotechnol.*, 7: 172-178.
- Nguyen, A. Q., Kim, Y. G., Kim, S. B. and Kim, C. J. (2018). Improved tolerance of recombinant *Escherichia coli* to the toxicity of crude glycerol by overexpressing trehalose biosynthetic genes (*otsBA*) for the production of b-carotene. *Biores. Technol.*, 143: 531-537.
- Okano, K., Tanaka, T., Ogino, C., Fukuda, H. and Kondo, A. (2019). Biotechnological production of enantiomeric pure lactic acid from renewable resources: recent achievements, perspectives and limits. *Appl. Microbiol. Biotechnol.*, 85: 413-423.
- Oshima, T. and Moriya, T. (2019). A preliminary analysis of microbial and biochemical properties of high-temperature compost. *Ann. NY Acad. Sci.*, 1125: 338-344.
- Peng, X., Qiao, W., Mi, S., Jia, X., Su, H. and Han, Y. (2019). Characterization of hemicellulase and cellulose from the extremely thermophilic bacterium *Caldicellulosiruptor owensensis* and their potential application for bioconversion of lignocellulosic biomass without pretreatment. *Biotechnol. Biofuels*, 8(131): 1-14.
- Rainey, F. A., Donnison, A. M., Jansen, P. H., Saul, D. and Rodrigo, A. (2019). Description of *Caldicellulosiruptor saccharolyticus* gen nov, sp nov: an obligately anaerobic, extremely thermophilic, cellulolytic bacterium. *FEMS Microbiol. Lett.*, 120: 263-266.



- Sahni, N. and Gupta, U. (2018). Bio-degradation of synthetic textile dyes by thermophilic lignolytic fungal isolates. *J. Adv. Lab. Res. Biol.* 5(4): 137-139.
- Santos, H. and Da Costa, M. S. (2016). Compatible solutes of organisms that live in hot saline environments. *Environ. Microbiol.*, 4: 501-509.
- Scully, S. M. and Orlygsson, J. (2019). Recent advances in second generation ethanol production by thermophilic bacteria. *Energies* 8(1): 1-30.
- Sommer, P., Georgieva, T. and Ahring, B. K. (2016). Potential for using thermophilic anaerobic bacteria for bioethanol production from hemicelluloses. *Biochem. Soc. Trans.*, 32(2): 283-289.
- Sprott, G. D., Meloche, M. and Richards, J. C. (2018). Proportions of diether, macrocyclic diether, and tetraether lipids in *Methanococcus jannaschii* grown at different temperatures. *J. Bacteriol.*, 173(12): 3907-3910.
- Thompson, M. J. and Eisenberg, D. (2018). Transproteomic evidence of a loop-deletion mechanism for enhancing protein thermostability. *J. Mol. Biol.*, 290(2): 595-604.
- Varankovich, N. V., Nickerson, M. T. and Korber, D. R. (2019). Probiotic-based strategies for therapeutic and prophylactic use against multiple gastrointestinal diseases. *Front. Biotechnol.*, 6: 1-14.
- Verma, A. and Shirkot, P. (2018). Purification and characterization of thermostable laccase from thermophilic *Geobacillus thermocatenulatus* MS5 and its applications in removal of textile dyes. *Scholars Acad. J. Biosci.*, 2(8): 479-485.
- Viljoen, J. A., Fred, E. B. and Peterson, W. H. (2018). the fermentation of cellulose by thermophilic bacteria. *J. Agric. Sci.*, 16: 1-17.
- Von Sivers, M. and Zacchi, G. (2020). A techno-economical comparison of three processes for the production of ethanol from wood. *Biores. Technol.*, 51: 43-52.
- Weimer, P. J. (2020). Thermophilic anaerobic fermentation of hemicellulose and hemicelluloses derived aldose sugars by *Thermoanaerobacter* strain B6A. *Archives Microbiol.*, 143: 130-136.
- Wiegel, J. and Ljungdahl, L. G. (2019). The importance of thermophilic bacteria in biotechnology. *Crit. Rev. Biotechnol.*, 3: 39-108.
- Woodward, J., Orr, M., Cordray, K. and Greenbaum, E. (2018). Biotechnology: enzymatic production of biohydrogen. *Nature*, 405:1014-1015.
- Ye, X., Honda, K., Sakai, T., Okano, K., Omasa, T., Hirota, R., Kuroda, A. and Ohtake, H. (2019). Synthetic metabolic engineering—a novel, simple technology for designing a chimeric metabolic pathway. *Microb. Cell Fact.*, 11: 120-131.